

**Title:** Trends in agricultural triazole fungicide use in the United States, 1992–2016 and possible implications for antifungal-resistant fungi in human disease

**Authors:** Mitsuru Toda, MS, PhD<sup>1</sup>, Karlyn D. Beer, MS, PhD<sup>1</sup>, Kathryn M. Kuivila, MS, PhD<sup>2</sup>, Tom M. Chiller, MD, MPHTM<sup>1</sup>, Brendan R. Jackson, MD, MPH<sup>1</sup>

**Affiliations:**

1. Mycotic Diseases Branch, Division of Foodborne, Waterborne, and Environmental Diseases (DFWED), National Center for Emerging and Zoonotic Infectious Diseases (NCEZID), Centers for Disease Control and Prevention (CDC)

2. U.S. Geological Survey, Oregon Water Science Center, Portland, Oregon

Corresponding author: Mitsuru Toda, [ [HYPERLINK "mailto:nrk7@cdc.gov" \h](mailto:nrk7@cdc.gov) ]; 1600 Clifton Road NE, Mailstop H24-9, Atlanta, Georgia 30329

**Competing financial interest:** The authors declare they have no actual or potential competing financial interests

**Abstract**

**Background**

The fungus *Aspergillus fumigatus* is the leading cause of invasive mold infections, which cause severe disease and death in immunocompromised people. Use of triazole antifungal medications in recent decades has

improved patient survival; however, triazole-resistant infections have become common in parts of Europe and are emerging in the United States. Triazoles are also a class of fungicides used in plant agriculture, and certain triazole-resistant *A. fumigatus* strains found causing disease in humans have been linked to environmental fungicide use.

## Objectives

We examined U.S. temporal and geographic trends in use of triazole fungicides using U.S. Geological Survey agricultural pesticide use estimates.

## Discussion

Based on our analysis, overall tonnage of triazole fungicide use nationwide was relatively constant during 1992–2005 but increased >4-fold during 2006–2016 to 2.9 million kg in 2016. During 1992–2005, triazole fungicide use occurred mostly in orchards and grapes, wheat, and other crops, but recent increases in use have occurred primarily in wheat, corn, soybeans, and other crops, particularly in Midwest and Southeast states. We conclude that given chemical similarities between triazole fungicides and triazole antifungal drugs used in human medicine, increased monitoring for environmental and clinical triazole resistance in *A. fumigatus* would improve overall understanding of these interactions, as well as help identify strategies to mitigate development and spread of resistance.

## Background

Invasive aspergillosis is a severe and frequently fatal fungal disease (mortality rate 25%–59%) that most commonly affects people who are immunocompromised (e.g., because of transplantation or malignancy) or have structural lung disease (e.g., chronic obstructive pulmonary disease (COPD)) (Kontoyiannis et al. 2010;

49 Pappas et al. 2010; Steinbach et al. 2012). Approximately 15,000 U.S. hospitalizations with invasive aspergillosis  
50 are estimated to occur annually based on medical coding data, with incidence increasing over the past decade,  
51 in part because of growing numbers of patients at risk (Benedict et al. 2019; Vallabhaneni et al. 2017). In high  
52 risk groups, such as solid organ transplantation recipients, incidence can approach 1% (Pappas et al. 2010).  
53 However, medical coding likely does not encompass all diagnosed cases, and the lack of national public health  
54 surveillance limits understanding of the true burden. Furthermore, many more undiagnosed cases likely exist. A  
55 systematic review of 31 studies of autopsy-confirmed misdiagnosis among intensive care unit patients during  
56 1966–2011 (5,863 examinations, 14 countries represented) indicated that aspergillosis was one of the most  
57 commonly missed diagnoses (Winters et al. 2012).

58  
59 *Aspergillus fumigatus*, the species of pathogenic fungi that causes most invasive aspergillosis (Patterson et al.  
60 2000), is common in the environment, particularly in decaying plant material but also at low levels in ambient air  
61 (Tekaia and Latgé 2005). Unlike many other fungi, it is thermotolerant up to 65 degrees Celsius and grows  
62 optimally at normal and febrile human body temperatures (roughly 37–40 degree Celsius), including during  
63 fever response, a key factor in its human pathogenicity, as well as at elevated temperatures found in composting  
64 organic matter (Kwon-Chung and Sugui 2013). Although it is widely present in agricultural areas, it is not known  
65 to cause disease in plants. Mold-active triazole antifungal medications (e.g., voriconazole) are the mainstay of  
66 treatment for invasive aspergillosis, having substantially improved patient survival following their introduction in  
67 the 1990s (Herbrecht et al. 2002; Verweij et al. 2016a). Only three main classes of antifungal medications  
68 (triazoles, echinocandins, and polyenes) are available to treat systemic fungal infections like aspergillosis.

69  
70 Whereas relatively few fungi cause invasive disease in humans, fungi are the most common cause of plant  
71 infections. Fungicides have been widely used for centuries to treat plant infections, prevent crop loss, and  
72 increase agricultural yield; fungicides are also used to preserve wood and other materials. (Morton and Staub

2008; Russell 2005; Kleinkauf and European Centre for Disease Prevention and Control 2013; US EPA 2015; Wise et al. 2019; Wise and Mueller 2011). Data on global triazole usage are limited, and the Food and Agriculture Organization provides data on combined triazole and diazole use, making it difficult to determine the amount of triazole use alone (FAOSTAT). Sales data suggest that triazoles are widely used agricultural fungicide classes, comprising over a quarter of estimated global fungicide sales (Kleinkauf and European Centre for Disease Prevention and Control 2013). Fungal pathogens of agricultural crops have developed resistance to many classes of fungicides, including triazoles (Cools and Fraaije 2008; Hu et al. 2016; Price et al. 2015), prompting the Fungicide Resistance Action Committee (FRAC) and other organizations to devote substantial resources to preventing and managing resistance (FRAC | Home). Notably, certain agricultural triazole fungicides, including bromuconazole, difenoconazole, epoxiconazole, propiconazole, and tebuconazole are structurally highly similar to medical triazoles used to treat aspergillosis (e.g., voriconazole, itraconazole, and posaconazole) (Snelders et al. 2012).

Like plant pathogens that have developed resistance to triazole fungicides, *A. fumigatus* strains resistant to medical triazoles have emerged globally, prompting public health concerns. Resistant aspergillosis is associated with treatment failure and high mortality, ranging from 42% to 88% (Lestrade et al. 2019; Resendiz-Sharpe et al. 2019; van der Linden et al. 2011). Death occurs more commonly in resistant infections, with 90-day mortality being 25% higher in patients with resistant versus susceptible aspergillosis in a European study (Lestrade et al. 2019). Resistance in *A. fumigatus* can develop in two ways. First, it can develop inside the body under selection pressure from long-term use of triazole medications. During the 1990s, small numbers of triazole-resistant infections were identified in patients receiving long-term triazole prophylaxis or therapy (e.g., for aspergilloma, cavitary lung disease, or other non-invasive aspergillosis), with resistance mechanisms involving point mutations in the triazole target and ergosterol synthesis gene, *CYP51A* (Camps et al. 2012; Heo et al. 2017; Howard et al. 2013, 2009). Resistance occurs less frequently in invasive aspergillosis, presumably because the fungus has less

time to grow in the body. Given the contribution of antifungal use to triazole resistance in *A. fumigatus*, it is notable that triazole use in U.S. hospitals declined by 21% during 2006–2012, the most recent years with available data (Vallabhaneni et al. 2018).

In the late 1990s, a new resistance mechanism was identified in patients who had *A. fumigatus* infections, and the same mechanism was identified in *A. fumigatus* exposed to triazole fungicides in the environment. This mechanism, TR34/L98H (which we will refer to as TR34), includes a 34-base pair tandem repeat (TR) in the *cyp51A* promoter coupled with a specific point mutation in the coding region and can confer resistance to all triazole medications, known as pan-resistance (Abdolrasouli et al. 2018). In contrast to the resistance mechanism that can develop inside the human body, this environmental resistance was observed in isolates primarily from patients who had never taken triazole medicines (Snelders et al. 2008; Verweij et al. 2007), with subsequent studies finding that 53%–64% of patients with resistant infection lacked exposure to medical triazoles (van der Linden et al. 2011, 2013).

Because triazoles are widely used in agriculture as fungicides, researchers suspected that the TR34-based resistance developed in the environment under fungicide-induced selection pressure (Bromley et al. 2014; Snelders et al. 2009) and that infections resulted from exposure to already-resistant *A. fumigatus* rather than resistance developing in the patient (Berger et al. 2017). Subsequent research provided additional evidence for this hypothesis and identified a second genotype, TR46/Y121F/T289A (TR46), thought to be linked to fungicide use (Astvad et al. 2014; Chowdhary et al. 2014b, 2015; Lavergne et al. 2015; Le Pape et al. 2016; Montesinos et al. 2014; Steinmann et al. 2015; van der Linden et al. 2013, 2015; Vermeulen et al. 2012). Although the TR-based mechanisms may not be definitive markers of environmental resistance, one report described a resistant isolate with a TR120 mechanism in a patient on long-term triazole therapy for chronic aspergillosis (Hare et al. 2019). Overall, evidence suggests that isolates with TR34 and TR46 mutations result from environmental triazole

121 exposure (Buil et al. 2019).

122

123 TR34 and TR46-mediated resistance has become common in patients with aspergillosis in parts of Europe, where

124 up to 20% of infections are now resistant to medical triazoles (Bueid et al. 2010; Lelièvre et al. 2013; Resendiz-

125 Sharpe et al. 2019; van der Linden et al. 2015; Vermeulen et al. 2013). Resistant *A. fumigatus* strains with TR34

126 and TR46 mutations have also been reported among azole-naïve patients in the Middle East, Asia, Africa,

127 Australia, and South America (Chowdhary et al. 2014a, 2017; Meis et al. 2016; Vermeulen et al. 2013; Verweij et

128 al. 2016a). In addition, environmental isolates with TR34 and TR46 mutations have been detected in Europe,

129 Asia, South America, and East Africa (Alvarez-Moreno et al. 2019; Badali et al. 2013; Chowdhary et al. 2012,

130 2014b; Dunne et al. 2017; Le Pape et al. 2016; Mortensen et al. 2010; Schoustra et al. 2019; Vermeulen et al.

131 2012). Further supporting a link between fungicide use and clinical resistance, triazole fungicides similar to

132 medical antifungals were introduced for agricultural use in the Netherlands just before the first TR34 strain was

133 found in human clinical settings in the late-1990s (Meis et al. 2016).

134

135 In the United States, associations between agricultural triazole fungicide use and human infections have not

136 been investigated, but a small number of infections caused by resistant *A. fumigatus* strains have been identified

137 (CDC 2019). The first TR-based resistance in patients was reported in 2016, including retrospectively identified

138 isolates (two TR<sub>34</sub> and two TR<sub>46</sub>) collected as early as 2008 (Vazquez and Manavathu 2016; Wiederhold et al.

139 2016). An additional 6 isolates were detected through 2018 (Beer 2018). Together, these 10 isolates likely reflect

140 only a small proportion of the true number of resistant infections given lack of standardized surveillance and

141 limited clinical testing. Resistant *A. fumigatus* strains with the TR<sub>34</sub> mutation have also been found in peanut crop

142 debris in the U.S. state of Georgia that had been treated with propiconazole and tebuconazole, triazoles that are

143 structurally similar to medical triazoles (Hurst et al. 2017), demonstrating this resistance was also present in the

144 U.S. agricultural environment. Because of this emergence in the United States, CDC has placed triazole-resistant

145 *A. fumigatus* on its “Watch List” for antimicrobial resistance threats (CDC 2019).

146

147 Given increased global incidence of triazole-resistant *Aspergillus* infections, recent identification of triazole  
148 resistance mechanisms linked to environmental agricultural fungicide use in the United States, and triazole  
149 agricultural fungicides with the same mechanism of action as triazole antifungal medications, we characterized  
150 trends in U.S. agricultural triazole use to explore possible implications for antifungal resistant human infections.  
151 We also examined available data regarding the use of triazole fungicides for purposes other than food  
152 production, including turf and other landscape maintenance and flower production.

153

## 154 **Methods**

155

156 We analyzed publicly available state-level estimates of annual agricultural pesticide use from the U.S. Geological  
157 Survey (USGS) (Baker and Stone 2015; Stone 2013; Thelin and Stone 2013) for 15 triazole fungicides used in the  
158 United States during 1992–2016 (USGS 2017). Data for District of Columbia, Hawaii, Alaska, and territories were  
159 not included in the estimates. Methods for these estimates are described in detail elsewhere (Baker and Stone  
160 2015; Stone 2013; Thelin and Stone 2013). Briefly, for states other than California, proprietary farm survey data  
161 collected by Gfk Kynetec, Inc. on amounts of pesticide use on specific crops are aggregated by the U.S.  
162 Department of Agriculture to estimate pesticide-by-crop use rates within Crop Reporting Districts (CRDs). Each  
163 CRD covers multiple counties and each county is assigned to a single CRD. County-level pesticide use estimates  
164 are then derived by applying CRD-level pesticide-by-crop use rates to county-level estimates of the harvested  
165 acreage of each relevant crop (based on U.S. Department of Agriculture’s Census of Agriculture data) and state-  
166 level use estimates are derived by summing the county-level estimates. When survey-based pesticide-by-crop  
167 use rates are missing for a CRD in a given year, two different approaches are used to account for the missing  
168 data (Thelin and Stone, 2013). Estimates based on the first approach assumes zero use for counties with missing

169 data and are referred to as “low” use estimates. Estimates based on the second approach extrapolates rates  
170 based on data for nearby CRDs and are referred to as “high” use estimates. Specifically, pesticide-by-crop use  
171 rates are estimated using the median rate for all contiguous CRDs; or, if data are missing for all contiguous CRDs,  
172 the median rate for all CRDs adjacent to contiguous CRDs; or, if data are missing for all of these CRDs, the  
173 median of all non-zero rates for all CRDs within the same USDA Farm Resource Region. To simplify  
174 interpretation, we used mean of the low and high annual agricultural pesticide use estimates in this report,  
175 rather than presenting each separately. For California, USGS inputs data on county-level pesticide use from the  
176 state’s Pesticide Use Reports (PUR), collected by the Department of Pesticide Regulation (California Department  
177 of Pesticide Regulation).

178  
179 Fifteen triazoles in the USGS dataset are used primarily as fungicides. Because seven of these triazoles  
180 (difenoconazole, metconazole, myclobutanil, propiconazole, prothiconazole, tebuconazole, and triadimefon)  
181 accounted for 93% of triazole use, we grouped the remaining eight fungicides (cyproconazole, fenbuconazole,  
182 flusilazole, flutriafol, ipconazole, tetraconazole, triadimenol, and triticonazole) into a single category. Three of  
183 the five agricultural triazoles documented to be structurally similar to medical triazoles (Snelders et al. 2012) are  
184 registered for use in the United States (difenoconazole, propiconazole, and tebuconazole).

185  
186 Based on USGS classifications, we grouped crops into eight categories: corn, cotton, orchards and grapes (stone  
187 fruit trees, citrus, nut trees, apples, pears, and grapevines), rice, soybeans, vegetables and fruit (vegetables and  
188 non-orchard fruit, including beans, peas, greens, berries, and melons), wheat, and other crops. The other crop  
189 category includes pasture and hay (cropland for pasture, fallow and idle cropland, pastureland, and other hay),  
190 alfalfa, sorghum, non-wheat grains, tobacco, peanuts, sugarcane, sugar beets, and other miscellaneous crops  
191 (Baker and Stone 2015; Stone 2013; Thelin and Stone 2013).

192



193 We characterized estimated U.S. triazole fungicide usage stratified by year, specific compounds, crop type, and  
194 geographical location. To aid in interpretation, we used mean of the low and high annual agricultural pesticide  
195 estimates rather than presenting each separately. We also examined state-specific use of triazoles, including by  
196 crop type, over 5 time periods (1992–1996, 1997–2001, 2002–2006, 2007–2011, and 2012–2016) and compared  
197 use during the periods 2012–2016 versus 1992–1996. To calculate differences over time, we summed the mean  
198 metric tons of fungicide use for years 2012–2016 and subtracted that value with mean metric tons for years  
199 1992–1996. All analysis was completed in R (Version 3.6.3, RStudio) and maps were created in ArcGIS (ArcGIS  
200 Desktop 10.5.1, Esri Inc.).

201  
202 Because triazole fungicides are used in the environment for purposes other than food production, we separately  
203 examined California's PUR data for 2017, the most recent year with available data, because the system includes  
204 data on wider range of uses than the USGS dataset (California Department of Pesticide Regulation 2017). We  
205 examined triazole use in turf (golf course turf, landscape maintenance, bermudagrass, rights of way, and  
206 turf/sod), ornamental (garland chrysanthemum, greenhouse flower, greenhouse plants in containers,  
207 greenhouse transplants, outdoor flower, outdoor plants in containers, and outdoor transplants), treated lumber,  
208 and other (airport, animal burrows, animal premise, beehive, Christmas tree, non-agricultural outdoor buildings,  
209 commercial storages or warehouses, commodity fumigation, dairy equipment, ditch bank, farm building,  
210 agricultural building, food processing plant, timberland forest, other fumigation, seed grass, greenhouse  
211 fumigation, household, industrial processing water, industrial site, industrial disposable water waste disposal  
212 systems, public health, regulatory pest control, research commodity, structural pest control.

213

## 214 **Results**

215

216 Estimated triazole fungicide use was relatively constant between 1992 (428 metric tons) and 2006 (539 metric

217 tons), but increased 434% from 2006 to 2016, to 2,880 metric tons (Figure 1, Table S1). Triazole use by  
218 compound differed over time (Figure 2A, Table S2). Estimated use of propiconazole and tebuconazole, the most  
219 widely used fungicides in 2016, increased little from 1992 to 2006, whereas use increased by 366% for  
220 propiconazole and 229% for tebuconazole during 2006–2016. First use of three newer triazoles difenoconazole,  
221 metconazole, and prothiconazole was reported after 2006, and usage increased to a total of 732 metric tons in  
222 2016. In contrast, estimated use of myclobutanil and triadimefon decreased during 1992–2016 (Figure 2A, Table  
223 S2).

224  
225 Estimated triazole fungicide use by crop type also changed substantially over time (Figure 2B, Table S3). During  
226 1992–2005, the primary use was on wheat, orchards and grapes, and other crops. Use on wheat began to  
227 increase markedly in 2007, with use increasing 683% during 2006–2016, resulting in the highest use amongst all  
228 crops in 2016 (1253 metric tons). Use on corn and soybeans also increased dramatically, with use on corn  
229 growing from 0 to 437 metric tons during 2006–2016, while use on soybeans increased from 61 to 361 metric  
230 tons. Use on other crops, rice, vegetables, and cotton increased steadily over time but at a slower rate. Use on  
231 orchards and grapes remained relatively constant (Figure 2B, Table S3).

232  
233 The estimated geographical distribution of triazole fungicide use shifted as use by crop type changed over time  
234 (Figure 3, Table S4, Table S5). The two states with the highest use during the 2012–2016 period, North Dakota  
235 (1,800 metric tons) and Georgia (1,008 metric tons), also had the largest increases since 1992–1996. This was  
236 primarily due to application on wheat in North Dakota and other crops, such as peanuts, in Georgia (Figure S1).  
237 Although California had the third highest usage during 2012–2016 (711 metric tons), application increased <50%  
238 since 1992–1996; triazoles were used primarily on orchards and grapes. The geographic shift is apparent as  
239 triazole use increased in the Midwest with wheat, corn, and soybeans (Figure S1, Table S4, Table S5).

240

241 In California, based on estimated PUR data in a single year, 5% of reported triazole fungicide use occurred in  
242 non-food production settings (e.g., turf, flowers, landscape maintenance) (Table S6).

243

## 244 Discussion

245

246 Based on our analysis of USGS estimates, overall U.S. triazole fungicide use in agriculture was relatively constant  
247 during 1992–2005 and increased >4-fold during 2006–2016 based on USGS estimates. Although estimated  
248 triazole usage increased in nearly every crop type and state over the period, the increase occurred primarily in  
249 wheat, corn, soybeans, and other crops in the Midwest and Southeast. These increases may have implications  
250 for triazole resistance in pathogenic fungi for humans, particularly in *A. fumigatus*, based on evidence from  
251 Europe and elsewhere (Bueid et al. 2010; Lelièvre et al. 2013; Resendiz-Sharpe et al. 2019). Given that resistance  
252 mutations previously associated with environmental triazole use have recently been detected in U.S. patient and  
253 environmental *A. fumigatus* isolates (Beer 2018; Hurst et al. 2017), additional study of the role of agricultural  
254 fungicides is warranted.

255

256 Several factors may explain the dramatic increase in U.S. triazole fungicide use after 2006, including increased  
257 corn production in response to higher prices, plant diseases in certain regions, ability to use new fungicides on  
258 field crops, and marketing of fungicides for use on field crops (Mueller et al. 2017; Wise and Mueller 2011). For  
259 example, when soybean rust caused by the fungus *Phakopsora pachyrhizi* was first identified in the United States  
260 in 2004, several fungicides were registered or granted emergency exemptions for treatment of soybeans,  
261 including myclobutanil, propiconazole, tebuconazole, and tetraconazole (Battaglin et al. 2011; Sconyers et al.  
262 2006; Wise and Mueller 2011). Another class of fungicides called strobilurins have been marketed to increase  
263 soybean and corn yield, frequently in combination with triazoles (Swoboda and Pedersen 2009; Wise and  
264 Mueller 2011). Fungicides are also used preemptively and in targeted ways in what are called insurance

265 applications, cover sprays, or prophylactic treatments when they are added to spray tanks being used to apply  
266 other pesticides like herbicides or insecticides (DiFonzo 2012). More research may be helpful to understand the  
267 reasons behind the large increases in triazole fungicides.

268

269 Because both triazoles and *A. fumigatus* can travel in the environment, exposure and resistance selection should  
270 be considered beyond the sites of application at agricultural fields. For example, triazoles have been detected in  
271 surface waters across the country (Battaglin et al. 2011; Nowell et al. 2018; Sanders et al. 2018; Smalling and  
272 Orlando 2011). Further, triazoles can be transported long distances in the atmosphere (Désert et al. 2018;  
273 Schummer et al. 2010), and residues have been detected in amphibians living in remote locations in the Sierra  
274 Nevada, dozens of miles downwind from where they were applied (Smalling et al. 2013). This mobility means  
275 that *A. fumigatus* in areas outside agricultural land may be exposed to triazoles, providing opportunity for  
276 resistance to develop. *A. fumigatus* spores, like spores of fungal plant pathogens, can travel long distances in the  
277 air (Brown and Hovmøller 2002). Triazole-resistant *A. fumigatus* isolates with fungicide-associated TR mutations  
278 have been found inside the homes and in the yards of aspergillosis patients, in hospital gardens, and in air  
279 samples taken from inside hospitals (Chowdhary et al. 2014b; Lavergne et al. 2017; van der Linden et al. 2013).

280

281 Data on non-food production uses of triazole fungicides in the United States were limited to a single state,  
282 California, where 5% of triazole fungicides in 2017 were used for turf, landscape, flowers, lumber, and other. This  
283 proportion is likely to be different in other states and nationally, and is an important topic of further study,  
284 particularly because some of these uses may be closer to population centers. Residential use of triazole  
285 fungicides could also be examined, since consumers can purchase some of these fungicides (e.g., propiconazole)  
286 in stores and online.

287

288 Important parallels can be drawn between challenges with agricultural use of medically important triazoles and

289 agricultural use of medically important antibacterial drugs. In recent years, the Food and Drug Administration  
290 has required that new antimicrobial drugs used in food-producing animals undergo a risk assessment to  
291 determine potential impacts on bacteria of human health concern (Center for Veterinary Medicine 2019a,  
292 2019b). Evaluation of potential human health impacts of agricultural triazole fungicide should be considered in  
293 more depth. Given that greater use of an antimicrobial is known to select for increased antimicrobial resistance,  
294 and that triazole-resistant infections are emerging in plants, greater triazole resistance in human pathogens may  
295 emerge as well (Chowdhary et al. 2013). Although detection of TR34 and TR46 has been limited in the United  
296 States to date (Beer 2018), surveillance, reporting, and susceptibility testing for *A. fumigatus* infections are not  
297 routinely conducted, suggesting that such infections are likely more widespread. For example, only 62% of the  
298 infectious disease doctors surveyed through the Emerging Infections Network in the United States reported  
299 having access to susceptibility testing for *A. fumigatus*, and such tests were not routinely ordered. Nevertheless,  
300 physicians reported seeing resistance in the United States, with 19% observing any triazole resistance and 7%  
301 pan-resistance. Fourteen percent were aware of a possible link to environmental fungicide use (Walker et al.  
302 2018). In contrast, testing for resistance in *A. fumigatus* in Europe is more widespread. The European Centre for  
303 Disease Prevention and Control recommends triazole antifungal susceptibility testing on all clinical *A. fumigatus*  
304 isolates when starting antifungal therapy (Kleinkauf and European Centre for Disease Prevention and Control  
305 2013).

306

307 Several limitations are inherent in this descriptive analysis of US fungicide use. First, the USGS data are estimates  
308 based on a proprietary farm survey (except for California, which has a state reporting system), and some degree  
309 of error is expected. In this descriptive analysis, we took the mean of the USGS low and high triazole estimates,  
310 which is a simplification involving differing estimates. Second, we did not adjust triazole usage by units of  
311 acreage treated, arable land by state or crop, restriction of certain crops in a state, and availability of seed  
312 treatment data, although these may be areas of further study. Finally, although available evidence points to

313 environmental fungicide use as a driver of TR-based triazole resistance in *A. fumigatus* globally, direct  
314 associations between quantity, use pattern, and timing of agricultural fungicide use and resistant human  
315 infections in the United States have not yet been established.

316

317 In the United States, research and partnerships may allow for opportunities to intervene early before *A.*  
318 *fumigatus* resistance becomes a larger clinical problem. First, more robust laboratory-based surveillance for *A.*  
319 *fumigatus* infections (Verweij et al. 2016b), including systematic antifungal susceptibility testing and microbiome  
320 studies, could better determine the burden of resistant infections, as well as geographic and temporal trends.  
321 Second, wider-scale environmental testing could assess the distribution of resistance in the environment and  
322 agricultural sector. Third, interdisciplinary One Health partnerships could identify ways to mitigate resistance,  
323 including exploring alternative fungicides and integrated pest management (Chowdhary et al. 2013; Fisher et al.  
324 2018). Finally, antifungal stewardship in human medicine plays an important role in judicious use of these  
325 limited and important medications (Fitzpatrick et al. 2020), and hospital stewardship programs have been shown  
326 to reduce the burden of antimicrobial-resistant human infections (Ananda-Rajah et al. 2012; Baur et al. 2017).  
327 These analyses demonstrate that triazole fungicide use in agriculture has increased >4-fold during 2006–2016 in  
328 the United States, driven primarily by increases in propiconazole and tebuconazole, with the largest increases in  
329 central parts of the United States. Exposure of *A. fumigatus* to fungicides can select for mutations that cause  
330 resistance to the primary antifungals used to treat human aspergillosis. Data on agricultural triazole use can  
331 inform further research, risk assessments, and policy decisions related to resistant fungal infections associated  
332 with patient illness and death.

333

#### 334 **Acknowledgments**

335

336 Work by K. Kuivila was funded by the U.S. Geological Survey Oregon Water Science Center. We would like to

337 thank Wesley Stone (U.S. Geological Survey) for his assistance with the annual agricultural pesticide use  
338 estimates, Erin Ricketts (former CDC epidemiology elective student) for her initial exploration on the topic, and  
339 Drs. Megin Nichols and Dawn Sievert (CDC) for helpful comments on the manuscript.

340

341 Disclaimer

342

343 The findings and conclusions in this report are those of the authors and do not necessarily represent the official  
344 position of the Centers for Disease Control and Prevention (CDC). Any use of trade, firm, or product names is for  
345 descriptive purposes only and does not imply endorsement by the U.S. Government.

## References

- Abdolrasouli A, Petrou MA, Park H, Rhodes JL, Rawson TM, Moore LSP, et al. 2018. Surveillance for Azole-Resistant *Aspergillus fumigatus* in a Centralized Diagnostic Mycology Service, London, United Kingdom, 1998–2017. *Front Microbiol* 9; doi:10.3389/fmicb.2018.02234.
- Alvarez-Moreno C, Lavergne R-A, Hagen F, Morio F, Meis JF, Le Pape P. 2019. Fungicide-driven alterations in azole-resistant *Aspergillus fumigatus* are related to vegetable crops in Colombia, South America. *Mycologia* 111:217–224; doi:10.1080/00275514.2018.1557796.
- Ananda-Rajah M, Slavin M, Thursky K. 2012. The case for antifungal stewardship. *Curr Opin Infect Dis* 25:107–115; doi:10.1097/QCO.0b013e32834e0680.
- Astvad KMT, Jensen RH, Hassan TM, Mathiasen EG, Thomsen GM, Pedersen UG, et al. 2014. First Detection of TR46/Y121F/T289A and TR34/L98H Alterations in *Aspergillus fumigatus* Isolates from Azole-Naive Patients in Denmark despite Negative Findings in the Environment. *Antimicrob Agents Chemother* 58:5096–5101; doi:10.1128/AAC.02855-14.
- Badali H, Vaezi A, Haghani I, Yazdanparast SA, Hedayati MT, Mousavi B, et al. 2013. Environmental study of azole-resistant *Aspergillus fumigatus* with TR34/L98H mutations in the *cyp51A* gene in Iran. *Mycoses* 56:659–663; doi:10.1111/myc.12089.
- Baker NT, Stone WW. 2015. Estimated Annual Agricultural Pesticide Use for Counties of the Conterminous United States, 2008–12. US Geol Surv Data Series 907: 9 p. [ [HYPERLINK](#) ]



370 "https://gcc02.safelinks.protection.outlook.com/?url=http%3A%2F%2Fdx.doi.org%2F10.3133%2Fds907&data=0  
371 4%7C01%7Ckkuivila%40usgs.gov%7Cd7d66408542f435b7a5f08d891a53c47%7C0693b5ba4b184d7b9341f32f40  
372 0a5494%7C0%7C0%7C637419486151522709%7CUnknown%7CTWFpbGZsb3d8eyJWljiMC4wLjAwMDAiLCJQIjoi  
373 V2luMzliLCJBTiI6IklhaWwiLCJXVCI6Mn0%3D%7C1000&sdata=jIXbphZg30Oe9vIV6crcV51pb%2Fz3UP0WRwdiBn  
374 cQcsA%3D&reserved=0" ]

375

376 Battaglin WA, Sandstrom MW, Kuivila KM, Kolpin DW, Meyer MT. 2011. Occurrence of Azoxystrobin,  
377 Propiconazole, and Selected Other Fungicides in US Streams, 2005–2006. *Water Air Soil Pollut* 218:307–322;  
378 doi:10.1007/s11270-010-0643-2.

379

380 Baur D, Gladstone BP, Burkert F, Carrara E, Foschi F, Döbele S, et al. 2017. Effect of antibiotic stewardship on the  
381 incidence of infection and colonisation with antibiotic-resistant bacteria and *Clostridium difficile* infection: a  
382 systematic review and meta-analysis. *Lancet Infect Dis* 17:990–1001; doi:10.1016/S1473-3099(17)30325-0.

383

384 Beer KD. 2018. Multidrug-Resistant *Aspergillus fumigatus* Carrying Mutations Linked to Environmental Fungicide  
385 Exposure — Three States, 2010–2017. *MMWR Morb Mortal Wkly Rep* 67; doi:10.15585/mmwr.mm6738a5.

386

387 Benedict K, Jackson BR, Chiller T, Beer KD. 2019. Estimation of Direct Healthcare Costs of Fungal Diseases in the  
388 United States. *Clin Infect Dis* 68:1791–1797; doi:10.1093/cid/ciy776.

389

390 Berger S, El Chazli Y, Babu AF, Coste AT. 2017. Azole Resistance in *Aspergillus fumigatus*: A Consequence of  
391 Antifungal Use in Agriculture? *Front Microbiol* 8; doi:10.3389/fmicb.2017.01024.

392

393 Bromley MJ, van Muijlwijk G, Fraczek MG, Robson G, Verweij PE, Denning DW, et al. 2014. Occurrence of azole-

394 resistant species of *Aspergillus* in the UK environment. *J Glob Antimicrob Resist* 2:276–279;  
395 doi:10.1016/j.jgar.2014.05.004.  
396  
397 Brown JKM, Hovmøller MS. 2002. Aerial Dispersal of Pathogens on the Global and Continental Scales and Its  
398 Impact on Plant Disease. *Science* 297:537–541; doi:10.1126/science.1072678.  
399  
400 Bueid A, Howard SJ, Moore CB, Richardson MD, Harrison E, Bowyer P, et al. 2010. Azole antifungal resistance in  
401 *Aspergillus fumigatus*: 2008 and 2009. *J Antimicrob Chemother* 65:2116–2118; doi:10.1093/jac/dkq279.  
402  
403 Buil JB, Hare RK, Zwaan BJ, Arendrup MC, Melchers WJG, Verweij PE. 2019. The fading boundaries between  
404 patient and environmental routes of triazole resistance selection in *Aspergillus fumigatus*. *PLOS Pathog*  
405 15:e1007858; doi:10.1371/journal.ppat.1007858.  
406  
407 Camps SMT, Linden JWM van der, Li Y, Kuijper EJ, Dissel JT van, Verweij PE, et al. 2012. Rapid Induction of  
408 Multiple Resistance Mechanisms in *Aspergillus fumigatus* during Azole Therapy: a Case Study and Review of the  
409 Literature. *Antimicrob Agents Chemother* 56:10–16; doi:10.1128/AAC.05088-11.  
410  
411 California Department of Pesticide Regulation. 2017 Annual Statewide Pesticide Use Report Indexed by  
412 Chemical. Available: [https://www.cdpr.ca.gov/docs/pur/pur17rep/statewide\\_ai\\_2017.htm](https://www.cdpr.ca.gov/docs/pur/pur17rep/statewide_ai_2017.htm) [accessed 18  
413 November 2020].  
414  
415 CDC. 2019. Antibiotic Resistance Threats in the United States, 2019. 148.  
416  
417 Center for Veterinary Medicine. 2019a. CVM GFI #152 Evaluating the Safety of Antimicrobial New Animal Drugs

418 with Regard to Their Microbiological Effects on Bacteria of Human Health Concern. US Food Drug Adm.  
419 Available: [ HYPERLINK "http://www.fda.gov/regulatory-information/search-fda-guidance-" \h ]documents/cvm-  
420 gfi-152-evaluating-safety-antimicrobial-new-animal-drugs-regard-their-microbiological-effects [accessed 5  
421 December 2019].  
422  
423 Center for Veterinary Medicine. 2019b. CVM GFI #209 The Judicious Use of Medically Important Antimicrobial  
424 Drugs in Food-Producing Animals. US Food Drug Adm. Available: [ HYPERLINK "http://www.fda.gov/regulatory-  
425 information/search-fda-guidance-documents/cvm-gfi-209-" \h ]judicious-use-medically-important-antimicrobial-  
426 drugs-food-producing-animals [accessed 5 December 2019].  
427  
428 Chowdhary A, Kathuria S, Xu J, Meis JF. 2013. Emergence of Azole-Resistant *Aspergillus fumigatus* Strains due to  
429 Agricultural Azole Use Creates an Increasing Threat to Human Health. *PLOS Pathog* 9:e1003633;  
430 doi:10.1371/journal.ppat.1003633.  
431  
432 Chowdhary A, Kathuria S, Xu J, Sharma C, Sundar G, Singh PK, et al. 2012. Clonal Expansion and Emergence of  
433 Environmental Multiple-Triazole-Resistant *Aspergillus fumigatus* Strains Carrying the TR34/L98H Mutations in  
434 the *cyp51A* Gene in India. *PLOS ONE* 7:e52871;  
435 doi:10.1371/journal.pone.0052871.  
436  
437 Chowdhary A, Sharma C, Hagen F, Meis JF. 2014a. Exploring azole antifungal drug resistance in *Aspergillus*  
438 *fumigatus* with special reference to resistance mechanisms. *Future Microbiol* 9:697–711;  
439 doi:10.2217/fmb.14.27.  
440  
441 Chowdhary A, Sharma C, Kathuria S, Hagen F, Meis JF. 2015. Prevalence and mechanism of triazole resistance in

442 *Aspergillus fumigatus* in a referral chest hospital in Delhi, India and an update of the situation in Asia. *Front*  
 443 *Microbiol* 6; doi:10.3389/fmicb.2015.00428.

444

445 Chowdhary A, Sharma C, Meis JF. 2017. Azole-Resistant Aspergillosis: Epidemiology, Molecular Mechanisms, and  
 446 Treatment. *J Infect Dis* 216:S436–S444; doi:10.1093/infdis/jix210.

447

448 Chowdhary A, Sharma C, van den Boom M, Yntema JB, Hagen F, Verweij PE, et al. 2014b. Multi-azole-resistant  
 449 *Aspergillus fumigatus* in the environment in Tanzania. *J Antimicrob Chemother* 69:2979–2983;  
 450 doi:10.1093/jac/dku259.

451

452 Cools HJ, Fraaije BA. 2008. Are azole fungicides losing ground against *Septoria* wheat disease? Resistance  
 453 mechanisms in *Mycosphaerella graminicola*. *Pest Manag Sci* 64:681–684; doi:10.1002/ps.1568.

454

455 Désert M, Ravier S, Gille G, Quinapallo A, Armengaud A, Pochet G, et al. 2018. Spatial and temporal distribution  
 456 of current-use pesticides in ambient air of Provence-Alpes-Côte-d’Azur Region and Corsica, France. *Atmos*  
 457 *Environ* 192:241–256; doi:10.1016/j.atmosenv.2018.08.054.

458

459 DiFonzo C. 2012. The hidden costs of insurance pesticide applications to field crops. MSU Ext. Available: [https://\[](https://www.canr.msu.edu/news/the_hidden_costs_of_insurance_pesticide_applications_to_field_crops)  
 460 [HYPERLINK "http://www.canr.msu.edu/news/the\\_hidden\\_costs\\_of\\_insurance\\_pesticide\\_applications\\_to\\_fi"](http://www.canr.msu.edu/news/the_hidden_costs_of_insurance_pesticide_applications_to_field_crops) \h  
 461 [ield\\_crops](http://www.canr.msu.edu/news/the_hidden_costs_of_insurance_pesticide_applications_to_field_crops) [accessed 23 October 2019].

462

463 Dunne K, Hagen F, Pomeroy N, Meis JF, Rogers TR. 2017. Intercountry Transfer of Triazole-Resistant *Aspergillus*  
 464 *fumigatus* on Plant Bulbs. *Clin Infect Dis* 65:147–149; doi:10.1093/cid/cix257.

465

[ ADDIN ZOTERO\_BIBL {"uncited":[],"omitted":[],"custom":[]} CSL\_BIBLIOGRAPHY ]

Fisher MC, Hawkins NJ, Sanglard D, Gurr SJ. 2018. Worldwide emergence of resistance to antifungal drugs challenges human health and food security. *Science* 360:739–742; doi:10.1126/science.aap7999.

Fitzpatrick MA, Albarillo F, Santarossa M, Evans CT, Suda KJ. 2020. Variability in antifungal stewardship strategies among Society for Healthcare Epidemiology of America (SHEA) Research Network facilities. *Infect Control Hosp Epidemiol* 41:585–589; doi:10.1017/ice.2020.76.

FRAC | Home. Available: <http://www.frac.info/home> [accessed 21 October 2019].

Hare RK, Gertsen JB, Astvad KMT, Degn KB, Løkke A, Stegger M, et al. 2019. In Vivo Selection of a Unique Tandem Repeat Mediated Azole Resistance Mechanism (TR120) in *Aspergillus fumigatus* cyp51A, Denmark - Volume 25, Number 3—March 2019 - *Emerging Infectious Diseases journal* -411 CDC.; doi:10.3201/eid2503.180297.

Heo ST, Tataru AM, Jiménez-Ortigosa C, Jiang Y, Lewis RE, Tarrand J, et al. 2017. Changes in In Vitro Susceptibility Patterns of *Aspergillus* to Triazoles and Correlation With Aspergillosis Outcome in a Tertiary Care Cancer Center, 1999-2015. *Clin Infect Dis Off Publ Infect Dis Soc Am* 65:216–225; doi:10.1093/cid/cix297.

Herbrecht R, Denning DW, Patterson TF, Bennett JE, Greene RE, Oestmann J-W, et al. 2002. Voriconazole versus amphotericin B for primary therapy of invasive aspergillosis. *N Engl J Med* 347:408–415; doi:10.1056/NEJMoa020191.

Howard SJ, Cerar D, Anderson MJ, Albarrag A, Fisher MC, Pasqualotto AC, et al. 2009. Frequency and Evolution of

490 Azole Resistance in *Aspergillus fumigatus* Associated with Treatment Failure. *Emerg Infect Dis* 15:1068–1076;  
 491 doi:10.3201/eid1507.090043.  
 492  
 493 Howard SJ, Pasqualotto AC, Anderson MJ, Leatherbarrow H, Albarrag AM, Harrison E, et al. 2013. Major  
 494 variations in *Aspergillus fumigatus* arising within aspergillomas in chronic pulmonary aspergillosis. *Mycoses*  
 495 56:434–441; doi:10.1111/myc.12047.  
 496  
 497 Hu M-J, Cox KD, Schnabel G. 2016. Resistance to Increasing Chemical Classes of Fungicides by Virtue of  
 498 “Selection by Association” in *Botrytis cinerea*. *Phytopathology*<sup>TM</sup> 106:1513–1520; doi:10.1094/PHYTO-04-16-  
 499 0161-R.  
 500  
 501 Hurst SF, Berkow EL, Stevenson KL, Litvintseva AP, Lockhart SR. 2017. Isolation of azole-resistant *Aspergillus*  
 502 *fumigatus* from the environment in the south-eastern USA. *J Antimicrob Chemother* 72:2443–2446;  
 503 doi:10.1093/jac/dkx168.  
 504  
 505 Kleinkauf N, European Centre for Disease Prevention and Control, eds. 2013. *Risk assessment on the impact of*  
 506 *environmental usage of triazoles on the development and spread of resistance to medical triazoles in Aspergillus*  
 507 *species*. ECDC [u.a.] Europäisches Zentrum für die Prävention und die Kontrolle von Krankheiten:Stockholm.  
 508  
 509 Kontoyiannis DP, Marr KA, Park BJ, Alexander BD, Anaissie EJ, Walsh TJ, et al. 2010. Prospective surveillance for  
 510 invasive fungal infections in hematopoietic stem cell transplant recipients, 2001-2006: overview of the  
 511 Transplant-Associated Infection Surveillance Network (TRANSNET) Database. *Clin Infect Dis Off Publ Infect Dis*  
 512 *Soc Am* 50:1091–1100; doi:10.1086/651263.  
 513

514 Kwon-Chung KJ, Sugui JA. 2013. *Aspergillus fumigatus*—What Makes the Species a Ubiquitous Human Fungal  
 515 Pathogen? PLoS Pathog 9; doi:10.1371/journal.ppat.1003743.

516

517 Lavergne R-A, Chouaki T, Hagen F, Toublanc B, Dupont H, Jounieaux V, et al. 2017. Home Environment as a  
 518 Source of Life-Threatening Azole-Resistant *Aspergillus fumigatus* in Immunocompromised Patients: Table 1. Clin  
 519 Infect Dis 64:76–78; doi:10.1093/cid/ciw664.

520

521 Lavergne R-A, Morio F, Favennec L, Dominique S, Meis JF, Gargala G, et al. 2015. First description of azole-  
 522 resistant *Aspergillus fumigatus* due to TR46/Y121F/T289A mutation in France. Antimicrob Agents Chemother  
 523 59:4331–4335; doi:10.1128/AAC.00127-15.

524

525 Le Pape P, Lavergne R-A, Morio F, Alvarez-Moreno C. 2016. Multiple Fungicide-Driven Alterations in Azole-  
 526 Resistant *Aspergillus fumigatus*, Colombia, 2015. Emerg Infect Dis 22:156–157; doi:10.3201/eid2201.150978.

527

528 Lelièvre L, Groh M, Angebault C, Maherault A-C, Didier E, Bougnoux M-E. 2013. Azole resistant *Aspergillus*  
 529 *fumigatus*: An emerging problem. Médecine Mal Infect 43:139–145; doi:10.1016/j.medmal.2013.02.010.

530

531 Lestrade PP, Bentvelsen RG, Schauwvlieghe AFAD, Schalekamp S, van der Velden WJFM, Kuiper EJ, et al. 2019.  
 532 Voriconazole Resistance and Mortality in Invasive Aspergillosis: A Multicenter Retrospective Cohort Study. Clin  
 533 Infect Dis 68:1463–1471; doi:10.1093/cid/ciy859.

534

535 Meis JF, Chowdhary A, Rhodes JL, Fisher MC, Verweij PE. 2016. Clinical implications of globally emerging azole  
 536 resistance in *Aspergillus fumigatus*. Philos Trans R Soc B Biol Sci 371:20150460; doi:10.1098/rstb.2015.0460.

537

538 Montesinos I, Dodemont M, Lagrou K, Jacobs F, Etienne I, Denis O. 2014. New case of azole-resistant *Aspergillus*  
 539 *fumigatus* due to TR46/Y121F/T289A mutation in Belgium. *J Antimicrob Chemother* 69:3439–3440;  
 540 doi:10.1093/jac/dku289.  
 541  
 542 Mortensen KL, Mellado E, Lass-Flörl C, Rodriguez-Tudela JL, Johansen HK, Arendrup MC. 2010. Environmental  
 543 Study of Azole-Resistant *Aspergillus fumigatus* and Other *Aspergilli* in Austria, Denmark, and Spain. *Antimicrob*  
 544 *Agents Chemother* 54:4545–4549; doi:10.1128/AAC.00692-10.  
 545  
 546 Morton V, Staub T. 2008. A Short History of Fungicides.  
 547  
 548 Mueller DS, Wise KA, Dufault NS, Bradley CA, Chilvers MI. 2017. *Fungicides for Field Crops | Mycology*. APS  
 549 Publications.  
 550  
 551 Nowell LH, Moran PW, Schmidt TS, Norman JE, Nakagaki N, Shoda ME, et al. 2018. Complex mixtures of  
 552 dissolved pesticides show potential aquatic toxicity in a synoptic study of Midwestern U.S. streams. *Sci Total*  
 553 *Environ* 613–614:1469–1488; doi:10.1016/j.scitotenv.2017.06.156.  
 554  
 555 Pappas PG, Alexander BD, Andes DR, Hadley S, Kauffman CA, Freifeld A, et al. 2010. Invasive Fungal Infections  
 556 among Organ Transplant Recipients: Results of the Transplant-Associated Infection Surveillance Network  
 557 (TRANSNET). *Clin Infect Dis* 50:1101–1111; doi:10.1086/651262.  
 558  
 559 Patterson TF, Kirkpatrick WR, White M, Hiemenz JW, Wingard JR, Dupont B, et al. 2000. Invasive aspergillosis.  
 560 Disease spectrum, treatment practices, and outcomes. I3 *Aspergillus* Study Group. *Medicine (Baltimore)* 79:250–  
 561 260; doi:10.1097/00005792-200007000-00006.



562

563 Price CL, Parker JE, Warrilow AGS, Kelly D, Kelly SL. 2015. Azole fungicides - understanding resistance  
564 mechanisms in agricultural fungal pathogens. *Pest Manag Sci* 71:1054–1058; doi:10.1002/ps.4029.

565

566 Resendiz-Sharpe A, Mercier T, Lestrade PPA, van der Beek MT, von dem Borne PA, Cornelissen JJ, et al. 2019.  
567 Prevalence of voriconazole-resistant invasive aspergillosis and its impact on mortality in haematology patients. *J*  
568 *Antimicrob Chemother* 74:2759–2766; doi:10.1093/jac/dkz258.

569

570 Russell PE. 2005. A century of fungicide evolution. *J Agric Sci* 143:11–25; doi:10.1017/S0021859605004971.

571

572 Sanders CJ, Orlando JL, Hladik ML. 2018. Detections of Current-Use Pesticides at 12 Surface Water Sites in  
573 California During a 2-Year Period Beginning in 2015. *US Geol Surv Data Series* 1088: 40 p.

574

575 Schoustra SE, Debets AJM, Rijs AJMM, Zhang J, Snelders E, Leendertse PC, et al. 2019. Environmental Hotspots  
576 for Azole Resistance Selection of *Aspergillus fumigatus*, the Netherlands. *Emerg Infect Dis* 25:1347–1353;  
577 doi:10.3201/eid2507.181625.

578

579 Schummer C, Mothiron E, Appenzeller BMR, Rizet A-L, Wennig R, Millet M. 2010. Temporal variations of  
580 concentrations of currently used pesticides in the atmosphere of Strasbourg, France. *Environ Pollut* 158:576–  
581 584; doi:10.1016/j.envpol.2009.08.019.

582

583 Sconyers LE, Kemeraite RC, Brock J, Phillips DV, Jost PH, Sikora EJ, et al. 2006. Asian Soybean Rust Development in  
584 2005: A Perspective from the Southeastern United States. Online. APSnet Features. doi:  
585 10.1094/APSnetFeatures-2006-0106

586

587 Smalling KL, Fellers GM, Kleeman PM, Kuivila KM. 2013. Accumulation of pesticides in pacific chorus frogs (  
588 *Pseudacris regilla*) from California's Sierra Nevada Mountains, USA: Pesticides residues in amphibians. Environ  
589 Toxicol Chem 32:2026–2034; doi:10.1002/etc.2308.

590

591 Smalling KL, Orlando JL. 2011. Occurrence of Pesticides in Surface Water and Sediments from Three Central  
592 California Coastal Watersheds, 2008–09. US Geol Surv Data Series 600: 70 p.

593

594 Snelders E, Camps SMT, Karawajczyk A, Schaftenaar G, Kema GHJ, Lee HA van der, et al. 2012. Triazole  
595 Fungicides Can Induce Cross-Resistance to Medical Triazoles in *Aspergillus fumigatus*. PLOS ONE 7:e31801;  
596 doi:10.1371/journal.pone.0031801.

597

598 Snelders E, Huis in 't Veld RAG, Rijs AJMM, Kema GHJ, Melchers WJG, Verweij PE. 2009. Possible Environmental  
599 Origin of Resistance of *Aspergillus fumigatus* to Medical Triazoles. Appl Environ Microbiol 75:4053–4057;  
600 doi:10.1128/AEM.00231-09.

601

602 Snelders E, Lee HAL van der, Kuijpers J, Rijs AJMM, Varga J, Samson RA, et al. 2008. Emergence of Azole  
603 Resistance in *Aspergillus fumigatus* and Spread of a Single Resistance Mechanism. PLOS Med 5:e219;  
604 doi:10.1371/journal.pmed.0050219.

605

606 Steinbach WJ, Marr KA, Anaissie EJ, Azie N, Quan S-P, Meier-Kriesche H-U, et al. 2012. Clinical epidemiology of  
607 960 patients with invasive aspergillosis from the PATH Alliance registry. J Infect 65:453–464;  
608 doi:10.1016/j.jinf.2012.08.003.

609

610 Steinmann J, Hamprecht A, Vehreschild MJGT, Cornely OA, Buchheidt D, Spiess B, et al. 2015. Emergence of  
 611 azole-resistant invasive aspergillosis in HSCT recipients in Germany. *J Antimicrob Chemother* 70:1522–1526;  
 612 doi:10.1093/jac/dku566.  
 613  
 614 Stone WW. 2013. Estimated Annual Agricultural Pesticide Use for Counties of the Conterminous United States,  
 615 1992–2009. Geol Surv Data Series 752. 1-p. pamphlet, 14 tables. [ HYPERLINK "<https://pubs.usgs.gov/ds/752/>" ]  
 616  
 617 Swoboda C, Pedersen P. 2009. Effect of Fungicide on Soybean Growth and Yield. *Agron J Madison* 101: 352–356.  
 618  
 619 Tejerina EE, Abril E, Padilla R, Ruíz CR, Ballen A, Frutos-Vivar F, et al. 2019. Invasive aspergillosis in critically ill  
 620 patients: An autopsy study. *Mycoses* 62:673–679; doi:10.1111/myc.12927.  
 621  
 622 Tejerina EE, Padilla R, Abril E, Frutos-Vivar F, Ballen A, Rodríguez-Barbero JM, et al. 2018. Autopsy-detected  
 623 diagnostic errors over time in the intensive care unit. *Hum Pathol* 76:85–90;  
 624 doi:10.1016/j.humpath.2018.02.025.  
 625  
 626 Tekaia F, Latgé J-P. 2005. *Aspergillus fumigatus*: saprophyte or pathogen? *Curr Opin Microbiol* 8:385–392;  
 627 doi:10.1016/j.mib.2005.06.017.  
 628  
 629 Thelin GP, Stone WW. 2013. Estimation of Annual Agricultural Pesticide Use for Counties of the Conterminous  
 630 United States, 1992–2009. U.S. Geological Survey Scientific Investigations Report 2013-5009, 54 p. [ HYPERLINK  
 631 "<http://pubs.usgs.gov/sir/2013/5009/>" ]  
 632  
 633 US EPA O. 2015. Overview of Wood Preservative Chemicals. US EPA. Available: [https://](https://www.epa.gov/pesticides/wood-preservative-chemicals)[ HYPERLINK

634 "http://www.epa.gov/ingredients-used-pesticide-products/overview-wood-preservative-" \h ]chemicals  
635 [accessed 1 May 2020].  
636  
637 USGS. 2017. USGS NAWQA: The Pesticide National Synthesis Project. Available:  
638 <https://water.usgs.gov/nawqa/pnsp/usage/maps/about.php> [accessed 20 July 2019].  
639  
640 Vallabhaneni S, Baggs J, Tsay S, Srinivasan AR, Jernigan JA, Jackson BR. 2018. Trends in antifungal use in US  
641 hospitals, 2006–12. *J Antimicrob Chemother* 73:2867–2875; doi:10.1093/jac/dky270.  
642  
643 Vallabhaneni S, Benedict K, Derado G, Mody RK. 2017. Trends in Hospitalizations Related to Invasive Aspergillosis  
644 and Mucormycosis in the United States, 2000–2013. *Open Forum Infect Dis* 4; doi:10.1093/ofid/ofw268.  
645  
646 van der Linden JWM, Arendrup MC, Warris A, Lagrou K, Pelloux H, Hauser PM, et al. 2015. Prospective  
647 Multicenter International Surveillance of Azole Resistance in *Aspergillus fumigatus*. *Emerg Infect Dis* 21:1041–  
648 1044; doi:10.3201/eid2106.140717.  
649  
650 van der Linden JWM, Camps SMT, Kampinga GA, Arends JPA, Debets-Ossenkopp YJ, Haas PJA, et al. 2013.  
651 Aspergillosis due to Voriconazole Highly Resistant *Aspergillus fumigatus* and Recovery of Genetically Related  
652 Resistant Isolates From Domiciles. *Clin Infect Dis* 57:513–520; doi:10.1093/cid/cit320.  
653  
654 van der Linden JWM, Snelders E, Kampinga GA, Rijnders BJA, Mattsson E, Debets-Ossenkopp YJ, et al. 2011.  
655 Clinical Implications of Azole Resistance in *Aspergillus fumigatus*, the Netherlands, 2007–2009 - Volume 17,  
656 Number 10—October 2011 - *Emerging Infectious Diseases journal* - CDC.; doi:10.3201/eid1710.110226.  
657

658 Vazquez JA, Manavathu EK. 2016. Molecular Characterization of a Voriconazole-Resistant, Posaconazole-  
659 Susceptible *Aspergillus fumigatus* Isolate in a Lung Transplant Recipient in the United States. *Antimicrob Agents*  
660 *Chemother* 60:1129–1133; doi:10.1128/AAC.01130-15.

661

662 Vermeulen E, Lagrou K, Verweij PE. 2013. Azole resistance in *Aspergillus fumigatus*: a growing public health  
663 concern. *Curr Opin Infect Dis* 26:493–500; doi:10.1097/QCO.0000000000000005.

664

665 Vermeulen E, Maertens J, Schoemans H, Lagrou K. 2012. Azole-resistant *Aspergillus fumigatus* due to  
666 TR46/Y121F/T289A mutation emerging in Belgium, July 2012. *Eurosurveillance* 17:20326;  
667 doi:10.2807/es.17.48.20326-en.

668

669 Verweij PE, Chowdhary A, Melchers WJG, Meis JF. 2016a. Azole Resistance in *Aspergillus fumigatus*: Can We  
670 Retain the Clinical Use of Mold-Active Antifungal Azoles? *Clin Infect Dis* 62:362–368; doi:10.1093/cid/civ885.

671

672 Verweij PE, Lestrade PPA, Melchers WJG, Meis JF. 2016b. Azole resistance surveillance in *Aspergillus fumigatus*:  
673 beneficial or biased? *J Antimicrob Chemother* 71:2079–2082; doi:10.1093/jac/dkw259.

674

675 Verweij PE, Mellado E, Melchers WJG. 2007. Multiple-Triazole-Resistant Aspergillosis. *N Engl J Med* 356:1481–  
676 1483; doi:10.1056/NEJMc061720.

677

678 Walker TA, Lockhart SR, Beekmann SE, Polgreen PM, Santibanez S, Mody RK, et al. 2018. Recognition of Azole-  
679 Resistant Aspergillosis by Physicians Specializing in Infectious Diseases, United States. *Emerg Infect Dis* 24:111–  
680 113; doi:10.3201/eid2401.170971.

681

682 Wiederhold NP, Gil VG, Gutierrez F, Lindner JR, Albataineh MT, McCarthy DJ, et al. 2016. First Detection of TR34  
683 L98H and TR46 Y121F T289A Cyp51 Mutations in *Aspergillus fumigatus* Isolates in the United States. *J Clin*  
684 *Microbiol* 54:168–171; doi:10.1128/JCM.02478-15.

685

686 Winters B, Custer J, Galvagno SM, Colantuoni E, Kapoor SG, Lee H, et al. 2012. Diagnostic errors in the intensive  
687 care unit: a systematic review of autopsy studies. *BMJ Qual Saf* 21:894–902; doi:10.1136/bmjqs-2012-000803.

688

689 Wise K, Mueller D. 2011. Are Fungicides No Longer Just For Fungi? An Analysis of Foliar Fungicide Use in Corn.  
690 *APSnet Feature Artic*; doi:10.1094/APSnetFeature-2011-0531.

691

692 Wise KA, Smith D, Freije A, Mueller DS, Kandel Y, Allen T, et al. 2019. Meta-analysis of yield response of foliar  
693 fungicide-treated hybrid corn in the United States and Ontario, Canada. *PLOS ONE* 14(6):e0217510;  
694 doi:10.1371/journal.pone.0217510.

695 Figure 1: Average agricultural triazole fungicide use by year in metric tons, United

696 States, 1992–2016

697

698 Estimates were derived by averaging “low” and “high” USGS agricultural pesticide estimates for each

699 year.

700

701 For corresponding numeric data, see Table S1.

702

703 Data from USGS. 2017. USGS NAWQA: The Pesticide National Synthesis Project.

EMBARGOED

704 Figure 2: Average agricultural triazole fungicide use by crop and compound type in  
705 metric tons, United States, 1992–2016

706

707 A. Triazole use by compound type in metric tons, 1992–  
708 2016

709

710 Fifteen triazoles included in the USGS dataset were grouped into 8 triazole categories:

711 1. Difenconazole

712 2. Metconazole

713 3. Myclobutanil

714 4. Other

715 5. Propiconazole

716 6. Prothiconazole

717 7. Tebuconazole

718 8. Triadimefon

719

720 The following triazoles were grouped into other triazole compound type category: cyproconazole,  
721 fenbuconazole, flusilazole, flutriafol, ipconazole, tetraconazole, triadimenol, and triticonazole.

722

723 For corresponding numeric data, see Table S2.

724

725 B. Triazole use by crop type in metric tons, 1992–2016

726

727 Crops were grouped into 8 categories:



728

729 1. Corn

730 2. Cotton

731 3. Orchards and grapes (stone fruit trees, citrus, nut trees, apples, pears, and grapevines)

732 4. Other crops

733 5. Rice

734 6. Soybeans

735 7. Vegetables and fruit (all vegetables and non-orchard fruit, including beans, peas, greens, berries,  
736 and melons)

737 8. Wheat

738

739 The following crop combinations were grouped into other crop type category: Pasture and Hay  
740 (cropland for pasture, fallow and idle cropland, pastureland, and other hay); Alfalfa; and Other  
741 (sorghum, non-wheat grains, tobacco, peanuts, sugarcane, sugar beets, and other miscellaneous crops).

742

743 For corresponding numeric data, see Table S3.

744

745 Data from USGS. 2017. USGS NAWQA: The Pesticide National Synthesis Project.

746

747 Estimates were derived by averaging “low” and “high” USGS agricultural pesticide estimates for each  
748 year.

749 Figure 3: Agricultural triazole fungicide usage map by state in metric tons, United States, 1992–2016

750

751 A: Differences in triazole fungicide usage 2012–16 and 1992–96 (in metric tons)

752

753 B: Triazole fungicide usage 1992–1996 (in metric tons)

754

755 C: Triazole fungicide usage 1997–2001 (in metric tons)

756

757 D: Triazole fungicide usage 2002–2006 (in metric tons)

758

759 E: Triazole fungicide usage 2007–2011 (in metric tons)

760

761 F: Triazole fungicide usage 2012–2016 (in metric tons)

762

763 Estimates from District of Columbia, Hawaii, Alaska, and the territories were not included in the maps.

764

765 For corresponding numeric data, see Table S4, S5, and S6.

766

767 Data from USGS. 2017. USGS NAWQA: The Pesticide National Synthesis Project.

768

769 Estimates were derived by averaging “low” and “high” USGS agricultural pesticide estimates for each

770 year.